

NEW SPECIES IN PHAEDRANASSA
AND EUCROSIA (AMARYLLIDACEAE)

Pierfelice Ravenna
INTA, Universidad de Chile, Santiago, Chile

New species of Amaryllidaceae, namely *Phaedranassa cinerea*, *Ph. viridilutea*, and *Eucrosia peruviana*, are described. The former two belong in the flora of Ecuador.

PHAEDRANASSA CINEREA Ravenna, sp. nov.

Bulbus late ovatus ad 9-11 cm longus et 7-8 cm latus inferne radicis crassis superne in pseudocollo 6-9 cm longo productus. Folia serotina post anthesin 2-3 cinereo-viridia utrinque pruinosa ad 45-51 cm longa et 7.5-10.5 mm lata; petioli erecto-patentes 15-16 cm longi inferne 9-11 mm lati in facie adaxiali subplani postice convexi, lamina oblanceolata patentia vel recurve patentia saepe undulata et plicatulo-nervosa acuta. Scapus teres glauco-viridis pruinosis ad 20-25 cm longus inferne leviter compressus circ. 9.7 mm latus ad apicem 5 mm latus. Spatha ad anthesin marcida valvis cartaceis lanceolatis ad basin liberis reflexis ad 33 mm longis. bracteae interiores 25-10 mm longae. Inflorescentia usque 11-flora. Pedicelli graciles glauco-virides teretifiliformes ad 27-30 mm longi. Flores declinati vel penduli. Ovarium ellipticum vel oblongo-ellipticum obtuso-triquetrum cinereo-viride pruinosum ad 7.7-8 mm longum circ. 3.5-3.8 mm latum. Perigonium leviter ventricosum circ. 31-35 mm longum. Perigonii tubus campanulatus vel poculiformis duis tertiiis inferioribus viridibus deinde carmineo-roseus. Tepala anguste oblanceolata usque 20 mm carmineo-rosea ad apicem oblique patentia cinereo-viridia albo-roseo-marginata; exteriora usque 29 mm longa in tertio inferiori 4 mm lata apicem versus 3.7-3.8 mm lata apiculo 1.8 mm longo intus ad basin tuberculo piloso notato; interiora usque 27 mm longa ad apicem 4.8-5 mm lata marginum albo-roseum 1.3 mm latum. Filamenta filiformi-subulata, sepalinum superiore 27.5 mm longum, sepalina lateralia 28.5 mm longa, petalinum inferiore 31 mm longum, petalina lateralia 32 mm longa. Antherae ellipticae versatiles sordidae 2.5-2.8 mm longae; pollen lutescens. Stylus filiformis ad initium anthesis 32 mm longus dein 38 mm longus. Stigma capitatus.

COLLECTIONS. Ecuador, prov. Chimborazo, road Gral. Eli-zalde (Bucay) - Pallatanga, 32 km from Pallatanga, 650 m; Øllgaard & Balslev 9006, 29-VIII-1976 (AAU, Herb. Rav.). Culta in Santiago Chilie ex bulbis a Dre. Øllgaard in Aequatore collectis; Ravenna 3049, XI-1982 (Herb. Ravenna holotype, AAU isotype).

Distinguishable by its large, ash green leaves, which are markedly pruinose on both faces. The flowers are shorter than in *Ph. dubia*.

PHAEDRANASSA VIRIDILUTEA Ravenna, sp. nov.

Planta usque 71 cm alta. Bulbus ovatus 40-45 mm longus et 35-40 mm latus in pseudocollo circ. 9 cm longo productus. Folia ad anthesin nulla. Scapus pariter cylindraceus usque 66 cm longus ad basin 7-8 mm latus et apicem 5 mm latus. Valvae lanceolatae marcescentes 40-42 mm longae; bracteae interiores anguste linearilanceolatae 30-20 mm longae. Pedicelli filiformes 11-30 mm longi. Flores patentes vel declinati. Ovarium 3.5-6.5 mm longum et 2.5-2.8 mm latum. Perigonii tubus viridis 9-10 mm longum et 2.5 mm latus. Perigonium anguste campanulatum vel subtubulare. Tepala oblanceolata circ. 11-12 mm lutea caeterum viride, exteriora 21-23 mm longa et 3.5-3.8 mm lata apiculo 0.6-0.7 mm longo, interiora 22-24 mm longa et 6.5-7 mm lata obtusa vel subacuta. Filamenta linearis attenuata, sepalina 15-16.5 mm longa, petalina 12-13 mm longae. Antherae ellipticae versatiles 2.5-3 mm longae. Stylus filiformis 3.5-3.7 mm longus. Stigma minute capitatus.

COLLECTIONS. Ecuador, prov. Azuay, highway Cuenca-Cola de San Pablo, km 54, 2580 m; Jef D. Boeke 973, 14-II-1977 (Herb. Ravenna holotype, NY isotype).

A distinct species by virtue of its flower colour.

EUCROSIA PERUVIANA Ravenna, sp. nov.

Planta 15-26 cm alta. Bulbus ovatus circ 20 mm latus tunicis exterioribus fusco-ochraceis: pseudocollus brevis. Folia serotina oblanceolato-petiolata fusco-viridia 11-16 cm longa et 13-25 mm lata; petiolus 46 mm longus. Scapus teres gracilis 13-25 cm longus. Valvae lanceolatae 4-10 mm longae. Inflorescentia 3-8-flora. Pedicelli 5-20 mm longi. Ovarium late ellipiticum vel ovatum obtuse triquetrum 3-4 mm longum et 2-2.5 mm latum. Perigonium bilabiate infundibulatum rubrum basin versus viride ad 16-22 mm longum et 12-17 mm in diametro horizontale et 16-24 mm in diametro verticali. Perigonii tubus viridis 3-4 mm longus. Tepala oblanceolata 14-17 mm longa extus ad apicem macula viridi: exteriora 4-5 mm lata apiculo 0.4-0.5 mm longo; interiora 5-7 mm lata obtusa vel subacuta. Filamenta fasciculata-declinata, sepalinum superius 50-60 mm longum, sepalina lateralia 54-65 mm longa, petalimum inferius 60-68 mm longum, petalina lateralia 65-70 mm longa. Antherae oblongo-reniformes luteae 2.5-3 mm longae. Stylus declinato-ascendens 65-80 mm longus. Stigma minute capitato-trilobatus.

COLLECTIONS. Perú, dept. Contumazá, prov. Guzmango, El Molino (Cascas—Contumazá; Sagástegui et al. 9338, 1-XI-1979 (TRP). Idem, Guzmango, 2500 m; López & Sagástegui, 5-X-1966 (TRP 6247). Idem, Cascas?, traído por un alumno; Sánchez-Vega 2790 (U. Tec. Cajamarca). Culta in Santiago ex bulbo in Cajamarca Peruviae collecto et a Dom. Sánchez-Vega donato; Ravenna 3070, XI-1983 (Herb. Ravenna, type).

Allied to *E. eucrosioides*, this species is separable by its more slender habit, narrower leaves, and a more regular perigone; the latter is green at the base, and uniformly red above.

ACKNOWLEDGEMENTS. The writer is indebted to Dr. B. Øllgaard, Botanical Institute, University of Aarhus, Denmark, and Prof. I. Sánchez Vega, Universidad Nacional Técnica de Cajamarca, Perú, for their consistent cooperation in sending dry and living material (bulbs) for study; Dr. A. López, and Prof. A. Sagástegui, Universidad de Trujillo, Perú, for facilities in the consultation of the Herbarium; and Dr. Patricia K. Holmgren, The New York Botanical Garden, for sending, among others, the specimen of *Phaedranassa viridilutea*.